

Jan Deleval

U.S. DEPARTMENT OF COMMERCE
Patent and Trademark Office

SEARCH REQUEST FORM

122380

Requestor's Name: Jennifer Kim Serial Number: 10/073,607
Date: 5/18/04 Phone: 20628 Art Unit: 1617

Search Topic:

Rem 4B-02

priority date 2/23/2000

Please write a detailed statement of search topic. Describe specifically as possible the subject matter to be searched. Define any terms that may have a special meaning. Give examples or relevant citations, authors, keywords, etc., if known. For sequences, please attach a copy of the sequence. You may include a copy of the broadest and/or most relevant claim(s).

1) Please search claims 31, 34, 35

2) Please search claims 9, 10, 12

RECEIVED
MAY 18 2004
610

THY

JW

STAFF USE ONLY

Date completed: 5/19/04
Searcher: JW
Terminal time: _____
Elapsed time: 15-00
CPU time: _____
Total time: _____
Number of Searches: _____
Number of Databases: _____

Search Site

☒ STIC
☐ CM-1
☐ Pre-S

Type of Search

☐ N.A. Sequence
☐ A.A. Sequence
☐ Structure
☒ Bibliographic

Vendors

☐ IG
☒ STN
☐ Dialog
☐ APS
☐ Geninfo
☐ SDC
☐ DARC/Questel
☐ Other



STIC Search Report

Biotech-Chem Library

STIC Database Tracking Number: 122380

TO: Jennifer Kim
Location: 4b02 / 4b18
Wednesday, May 19, 2004
Art Unit: 1617
Phone: 272-0628
Serial Number: 10 / 073607

From: Jan Delaval
Location: Biotech-Chem Library
Rem 1A51
Phone: 272-2504

jan.delaval@uspto.gov

Search Notes

=> d his

(FILE 'HOME' ENTERED AT 15:44:02 ON 19 MAY 2004)
SET COST OFF

FILE 'REGISTRY' ENTERED AT 15:44:21 ON 19 MAY 2004

E THIAZOLIDINEDIONE/CN
L1 1 S E3
E TROGLITAZONE/CN
L2 1 S E3
E CIGLITAZONE/CN
L3 1 S E3
E PIOGLITAZONE/CN
L4 1 S E3
E ROSIGLITAZONE/CN
L5 1 S E3
E ENGLITAZONE/CN
L6 1 S E3
E D-CHIRO-INOSITOL/CN
L7 1 S E3
E METFORMIN/CN
L8 2 S E3,E5
E CYTOPROTERONE/CN
E CITOPROTERONE/CN
E CYPROTERONE/CN
L9 2 S E3-E6
E FLUTAMIDE/CN
L10 1 S E3
E BICALUTAMIDE/CN
L11 1 S E3
E NILUTAMIDE/CN
L12 1 S E3
E RU-58841/CN
E RU 58841/CN
L13 1 S E3
E CANRENONE/CN
L14 1 S E3
E SPIRONOLACTONE/CN
L15 1 S E3
E PROGESTERONE/CN
L16 1 S E3
E 4MA/CN
E 4 MA/CN
E 4-MA/CN
L17 1 S E3
E KETOCONAZOLE/CN
L18 1 S E3
E CIMETIDINE/CN
L19 1 S E3
L20 9 S L1-L8
SEL RN
L21 227 S E1-E9/CRN
L22 217 S L21 NOT MXS/CI
L23 62 S L22 NOT (COMPD OR WITH)
L24 57 S L23 NOT IDS/CI
L25 12 S L9-L19
SEL RN
L26 284 S E10-E21/CRN
L27 17 S L26 NOT ((MXS OR IDS)/CI OR COMPD OR WITH OR UNSPECIFIED)
L28 15 S L27 NOT CONJUGATE

FILE 'HCAPLUS' ENTERED AT 15:58:46 ON 19 MAY 2004

L29 4690 S L20 OR L24

L30 2291 S D() (CHIROINOSITOL OR CHIRO INOSITOL) OR METFORMIN# OR DIMETHY
 L31 2672 S ROSIGLITAZON# OR BRL49653 OR BRL() (49653 OR 49 653) OR PIOGLI
 L32 2939 S CS045 OR CS 045 OR GR92132# OR GR() (92132# OR 92 132#) OR ADD
 L33 6815 S L29-L32
 L34 1970 S ISIS OR INSULIN(L) SENSITIV? (L) INCREAS? (L) SUBSTANC?
 L35 12995 S ALOPEC? OR BALDNESS OR BALD OR BALDING OR HAIR(L) (LOSS OR LOS
 L36 170 S PILOSEBAC?
 L37 3598 S HAIR(L) ?FOLLIC?
 L38 3520 S SCALP?
 L39 14 S L33 AND L35-L38
 L40 3 S L34 AND L35-L38
 L41 16 S L39,L40
 E HAIR/CT
 L42 16601 S E3-E18
 E E3+ALL
 L43 30975 S E6,E5+NT
 E E15+ALL
 L44 2450 S E13,E12+NT
 E E15+ALL
 E E17+ALL
 L45 20263 S E2+NT
 E E8+ALL
 E E19+ALL
 L46 2865 S E7,E6+NT
 E E16+ALL
 E E20+ALL
 L47 871 S E4
 E E6+ALL
 E E21+ALL
 L48 2215 S E3,E2+NT
 E HAIR/CT
 L49 65 S E86-E88
 L50 15265 S E44-E68
 L51 32 S L33 AND L42-L50
 L52 1 S L34 AND L42-L50
 L53 35 S L41,L51,L52
 L54 55669 S L25 OR L28
 L55 82367 S CYPROTERON? OR CYPROTERON?(S) ACETATE OR FLUTAMID? OR BICALUTA
 L56 213 S L55 AND L33
 L57 6 S L55 AND L34
 L58 14 S L56,L57 AND L53
 L59 764 S 17 BETA HYDROXY STEROID DEHYDROGENASE
 L60 1313 S 17 BETA HYDROXYSTEROID DEHYDROGENASE
 L61 320 S 3 ALPHA HYDROXY STEROID DEHYDROGENASE
 L62 736 S 3 ALPHA HYDROXYSTEROID DEHYDROGENASE
 L63 3272 S 5 ALPHA REDUCTASE

FILE 'REGISTRY' ENTERED AT 16:13:59 ON 19 MAY 2004

L64 3 S 9015-81-0 OR 9028-56-2 OR 9081-34-9

FILE 'HCAPLUS' ENTERED AT 16:14:07 ON 19 MAY 2004

L65 5023 S L64
 L66 6 S L33 AND L59-L63,L65
 L67 1 S L34 AND L59-L63,L65
 L68 6 S L66,L67
 L69 19 S L58,L68
 L70 4 S L69 AND ALOPEC?
 L71 4 S L35 AND L69
 L72 4 S L70,L71
 L73 175 S L54 AND L33,L34
 L74 3 S L73 AND L35-L38
 L75 11 S L73 AND L42-L50
 L76 19 S L69-L72,L74,L75

E KRAJCIK R/AU
L77 8 S E4,E6,E7
E ORENTREICH N/AU
L78 45 S E3,E4
L79 1 S US20020143039/PN OR (WO2001-US5653 OR US2000-184398#)/AP,PRN
L80 12 S L77-L79 AND L29-L63,L65
L81 1 S L80 AND L76
L82 11 S L80 NOT L81
L83 18 S L76 NOT L81
L84 6 S L83 AND (PD<=20000223 OR PRD<=20000223 OR AD<=20000223)
L85 12 S L83 NOT L84
SEL DN AN 1
L86 1 S L85 AND E1-E3
L87 2 S L79,L86
L88 2 S L87 AND L1-L28
L89 2 S L88 AND L29-L63,L65-L88

FILE 'REGISTRY' ENTERED AT 16:28:47 ON 19 MAY 2004

L90 1 S 56-03-1
L91 1 S 102-02-3

FILE 'HCAPLUS' ENTERED AT 16:28:58 ON 19 MAY 2004

L92 1269 S L90 OR L91
L93 4 S L92 AND L35-L38
L94 8 S L92 AND L42-L50
L95 8 S L93,L94
SEL DN AN 3-7
L96 5 S L95 AND E4-E18
L97 6 S L89,L96

=> fil hcaplus

FILE 'HCAPLUS' ENTERED AT 16:31:26 ON 19 MAY 2004

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2004 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 19 May 2004 VOL 140 ISS 21
FILE LAST UPDATED: 18 May 2004 (20040518/ED)

This file contains CAS Registry Numbers for easy and accurate substance identification.


=> d l97 all hitstr tot

L97 ANSWER 1 OF 6 HCAPLUS COPYRIGHT 2004 ACS on STN
AN 2004:271166 HCAPLUS
DN 140:292208
ED Entered STN: 02 Apr 2004
TI Use of a heterocyclic compound or one of its salts to stimulate or induce the growth of hair and/or to slow down its loss

IN Boulle, Christophe; Rozot, Roger; Dalko, Maria
 PA L'oreal, Fr.
 SO Fr. Demande, 32 pp.
 CODEN: FRXXBL
 DT Patent
 LA French
 IC ICM A61K007-075
 CC 62-3 (Essential Oils and Cosmetics)
 Section cross-reference(s): 28

FAN.CNT 2

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	FR 2845000	A1	20040402	FR 2002-12018	20020927
	WO 2004028441	A2	20040408	WO 2003-FR2823	20030925
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
PRAI	FR 2002-12018	A	20020927		
	US 2002-415462P	P	20021003		
OS	MARPAT 140:292208				
GI					



AB The invention refers to the use of a heterocyclic compound I or one of its salts in a capillary composition where Hy is a heterocyclic group with 4, 5, 6 or 7 atoms comprising possibly at least a carbonyl and/or a thiocarbonyl, the heterocyclic group being possibly substituted by at least a substituent chosen among a halogen, OR, SR, NRR', COR, CSR, NRCONR'R'', C(=NR)RR'', C(=NR)NR'R'', NRC(=NR')NR''R''', OCOR, COSR, SCOR, CSNRR', NRCSR', NRCSNR'R'', COOR, CONRR', CF3, CN, NRCOR', SO2R', SO2NRR', NRSO2R', saturated or unsatd. C1-20 alkyl radicals, cyclic group having at least a heteroatom, the alkyl radicals and cyclic group can be substituted, where R, R', R'', and R''', identical or different, are a hydrogen, a C1-20 alkyl radical or a substituted aryl radical; G represents O, S, NH; R1, R2 and R3 represent a hydrogen, a halogen, OR0, SR0, NR0R0', COR0, CSR0, NR0CONR0'R0'', C(=NR0)R0', C(=NR0)NR0'R0'', NR0C(=NR0')NR0''R0'', OCOR0, COSR0, SCOR0, CSNR0R0', NR0CSR0', NR0CSNR0'R0'', COOR0, CONR0R0', CF3, CN, NR0COR'0, SO2R0', SO2NR0R0', NR0SO2R0', a saturated or unsatd. C1-20 alkyl radical, at least a saturated or unsatd. cyclic compound containing at least a heteroatom, as **hair growth** stimulant agent. Thus, 4-{5-[(2,4-dioxo-1,3-thiazolidin-5-

ylidene)methyl]-2-furyl}benzoic acid (II) was prepared by the reaction of a furaldehyde phenylcarboxylic acid with thiozolidin-2,4-dione. A hair lotion contained II 0.80, propylene glycol 10.00, and iso-Pr alc. q.s. 100.00 g.

- ST heterocyclic compd salt **hair growth** stimulant
- IT Amino acids, biological studies
RL: COS (Cosmetic use); BIOL (Biological study); USES (Uses)
(N-acyl; use of heterocyclic compound or one of its salts to stimulate or induce **growth** of **hair** and/or to slow down its **loss**)
- IT Estrogens
RL: BSU (Biological study, unclassified); BIOL (Biological study)
(agonists and antagonists; use of heterocyclic compound or one of its salts to stimulate or induce **growth** of **hair** and/or to slow down its **loss**)
- IT Potassium channel
RL: COS (Cosmetic use); BIOL (Biological study); USES (Uses)
(agonists; use of heterocyclic compound or one of its salts to stimulate or induce **growth** of **hair** and/or to slow down its **loss**)
- IT Quaternary ammonium compounds, biological studies
RL: COS (Cosmetic use); BIOL (Biological study); USES (Uses)
(alkylbenzyl dimethyl, chlorides; use of heterocyclic compound or one of its salts to stimulate or induce **growth** of **hair** and/or to slow down its **loss**)
- IT Androgens
RL: COS (Cosmetic use); BIOL (Biological study); USES (Uses)
(antiandrogens; use of heterocyclic compound or one of its salts to stimulate or induce **growth** of **hair** and/or to slow down its **loss**)
- IT Ion channel blockers
(calcium; use of heterocyclic compound or one of its salts to stimulate or induce **growth** of **hair** and/or to slow down its **loss**)
- IT Tocopherols
RL: COS (Cosmetic use); BIOL (Biological study); USES (Uses)
(derivs.; use of heterocyclic compound or one of its salts to stimulate or induce **growth** of **hair** and/or to slow down its **loss**)
- IT Fatty acids, biological studies
RL: COS (Cosmetic use); BIOL (Biological study); USES (Uses)
(essential; use of heterocyclic compound or one of its salts to stimulate or induce **growth** of **hair** and/or to slow down its **loss**)
- IT Carboxylic acids, biological studies
RL: COS (Cosmetic use); BIOL (Biological study); USES (Uses)
(esters; use of heterocyclic compound or one of its salts to stimulate or induce **growth** of **hair** and/or to slow down its **loss**)
- IT Hair preparations
(**growth stimulants**; use of heterocyclic compound or one of its salts to stimulate or induce **growth** of **hair** and/or to slow down its **loss**)
- IT Carboxylic acids, biological studies
RL: COS (Cosmetic use); BIOL (Biological study); USES (Uses)
(hydroxy; use of heterocyclic compound or one of its salts to stimulate or induce **growth** of **hair** and/or to slow down its **loss**)
- IT Pruritus
(inhibitors; use of heterocyclic compound or one of its salts to stimulate or induce **growth** of **hair** and/or to slow down its **loss**)
- IT Prostaglandins

Steroids, biological studies

Tumor necrosis factors

RL: BSU (Biological study, unclassified); BIOL (Biological study)
(inhibitors; use of heterocyclic compound or one of its salts to
stimulate or induce **growth of hair** and/or to slow
down its **loss**)

IT **Hair preparations**
(lotions; use of heterocyclic compound or one of its salts to
stimulate or induce **growth of hair** and/or to slow
down its **loss**)

IT Essential oils

RL: COS (Cosmetic use); BIOL (Biological study); USES (Uses)
(peppermint; use of heterocyclic compound or one of its salts to
stimulate or induce **growth of hair** and/or to slow
down its **loss**)

IT Alcohols, biological studies

RL: COS (Cosmetic use); BIOL (Biological study); USES (Uses)
(polyhydric; use of heterocyclic compound or one of its salts to
stimulate or induce **growth of hair** and/or to slow
down its **loss**)

IT Interleukin 10

RL: COS (Cosmetic use); BIOL (Biological study); USES (Uses)
(promoters; use of heterocyclic compound or one of its salts to stimulate
or induce **growth of hair** and/or to slow down its
loss)

IT **Alopecia**

Antihistamines

Antimicrobial agents

Fungicides

Parasitocides

Vasodilators

(use of heterocyclic compound or one of its salts to stimulate or induce
growth of hair and/or to slow down its **loss**
)

IT Cytokines

Prostanoid receptors

RL: BSU (Biological study, unclassified); BIOL (Biological study)
(use of heterocyclic compound or one of its salts to stimulate or induce
growth of hair and/or to slow down its **loss**
)

IT Amino acids, biological studies

RL: COS (Cosmetic use); BIOL (Biological study); USES (Uses)
(use of heterocyclic compound or one of its salts to stimulate or induce
growth of hair and/or to slow down its **loss**
)

IT Carbohydrates, biological studies

RL: COS (Cosmetic use); BIOL (Biological study); USES (Uses)
(use of heterocyclic compound or one of its salts to stimulate or induce
growth of hair and/or to slow down its **loss**
)

IT Ceramides

RL: COS (Cosmetic use); BIOL (Biological study); USES (Uses)
(use of heterocyclic compound or one of its salts to stimulate or induce
growth of hair and/or to slow down its **loss**
)

IT Essential oils

RL: COS (Cosmetic use); BIOL (Biological study); USES (Uses)
(use of heterocyclic compound or one of its salts to stimulate or induce
growth of hair and/or to slow down its **loss**
)

IT Flavonoids

RL: COS (Cosmetic use); BIOL (Biological study); USES (Uses)
(use of heterocyclic compound or one of its salts to stimulate or induce
growth of hair and/or to slow down its **loss**)

- growth of hair and/or to slow down its loss
)
- IT Growth factors, animal
RL: COS (Cosmetic use); BIOL (Biological study); USES (Uses)
(use of heterocyclic compound or one of its salts to stimulate or induce
growth of hair and/or to slow down its loss
)
- IT Hormones, animal, biological studies
RL: COS (Cosmetic use); BIOL (Biological study); USES (Uses)
(use of heterocyclic compound or one of its salts to stimulate or induce
growth of hair and/or to slow down its loss
)
- IT Interleukin 1
RL: COS (Cosmetic use); BIOL (Biological study); USES (Uses)
(use of heterocyclic compound or one of its salts to stimulate or induce
growth of hair and/or to slow down its loss
)
- IT Interleukin 6
RL: COS (Cosmetic use); BIOL (Biological study); USES (Uses)
(use of heterocyclic compound or one of its salts to stimulate or induce
growth of hair and/or to slow down its loss
)
- IT Phospholipids, biological studies
RL: COS (Cosmetic use); BIOL (Biological study); USES (Uses)
(use of heterocyclic compound or one of its salts to stimulate or induce
growth of hair and/or to slow down its loss
)
- IT Protein hydrolyzates
RL: COS (Cosmetic use); BIOL (Biological study); USES (Uses)
(use of heterocyclic compound or one of its salts to stimulate or induce
growth of hair and/or to slow down its loss
)
- IT Proteins
RL: COS (Cosmetic use); BIOL (Biological study); USES (Uses)
(use of heterocyclic compound or one of its salts to stimulate or induce
growth of hair and/or to slow down its loss
)
- IT Retinoids
RL: COS (Cosmetic use); BIOL (Biological study); USES (Uses)
(use of heterocyclic compound or one of its salts to stimulate or induce
growth of hair and/or to slow down its loss
)
- IT Saponins
RL: COS (Cosmetic use); BIOL (Biological study); USES (Uses)
(use of heterocyclic compound or one of its salts to stimulate or induce
growth of hair and/or to slow down its loss
)
- IT Triterpenes
RL: COS (Cosmetic use); BIOL (Biological study); USES (Uses)
(use of heterocyclic compound or one of its salts to stimulate or induce
growth of hair and/or to slow down its loss
)
- IT Vitamins
RL: COS (Cosmetic use); BIOL (Biological study); USES (Uses)
(use of heterocyclic compound or one of its salts to stimulate or induce
growth of hair and/or to slow down its loss
)
- IT 141436-78-4, Protein kinase C
RL: BSU (Biological study, unclassified); BIOL (Biological study)
(activators; use of heterocyclic compound or one of its salts to
stimulate or induce growth of hair and/or to slow
down its loss)
- IT 58-82-2, Bradykinin 9032-92-2, Glycosidase 37255-34-8, Testosterone

5 α -**reductase** 63551-74-6, Arachidonate
lipoygenase 79955-99-0, Proteoglycanase 109300-99-4,
Glycosaminoglycanase

RL: BSU (Biological study, unclassified); BIOL (Biological study)
(inhibitors; use of heterocyclic compound or one of its salts to
stimulate or induce **growth of hair** and/or to slow
down its **loss**)

IT 50-28-2, Estradiol, biological studies 52-90-4, Cystein, biological
studies 57-13-6, Urea, biological studies 57-88-5, Cholesterol,
biological studies 59-67-6, Nicotinic acid, biological studies
59-67-6D, Nicotinic acid, esters 63-68-3, Methionine, biological studies
68-26-8D, Retinol, derivs. 69-72-7, Salicylic acid, biological studies
69-72-7D, Salicylic acid, derivs. 81-13-0, Panthenol 89-78-1, Menthol
97-59-6, Allantoin 98-79-3D, Pyroglutamic acid, esters 108-46-3,
Resorcinol, biological studies 108-95-2, Phenol, biological studies
113-92-8, Chlorpheniramine maleate 119-61-9, Benzophenone, biological
studies 121-54-0, Benzethonium chloride 137-08-6 302-79-4, Retinoic
acid 461-72-3, Hydantoin 526-95-4, Glycogenic acid 526-95-4D,
Glycogenic acid, acyl derivs. 19660-77-6D, Chlorophyllin, derivs.
38304-91-5, Minoxidil 68890-66-4, Octopirox 78418-01-6,
N-Octanoyl-5-salicylic acid 79217-60-0, Cyclosporin 104987-12-4D,
Ascomycin, derivs. 139615-39-7 675582-79-3 675582-81-7 675582-84-0
RL: COS (Cosmetic use); BIOL (Biological study); USES (Uses)

(use of heterocyclic compound or one of its salts to stimulate or induce
growth of hair and/or to slow down its **loss**
)

IT 331652-43-8P 357155-26-1P 675582-78-2P
RL: COS (Cosmetic use); SPN (Synthetic preparation); BIOL (Biological
study); PREP (Preparation); USES (Uses)
(use of heterocyclic compound or one of its salts to stimulate or induce
growth of hair and/or to slow down its **loss**
)

IT 1899-24-7, 5-Bromo-2-furaldehyde 2295-31-0, 2,4-
Thiazolidinedione 14047-29-1, 4-Carboxyphenylboronic acid
25487-66-5, 3-Carboxyphenylboronic acid 256658-04-5
RL: RCT (Reactant); RACT (Reactant or reagent)
(use of heterocyclic compound or one of its salts to stimulate or induce
growth of hair and/or to slow down its **loss**
)

IT 39245-15-3P 304884-54-6P
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT
(Reactant or reagent)
(use of heterocyclic compound or one of its salts to stimulate or induce
growth of hair and/or to slow down its **loss**
)

RE.CNT 11 THERE ARE 11 CITED REFERENCES AVAILABLE FOR THIS RECORD


RE

- (1) Anon; PATENT ABSTRACTS OF JAPAN 1998, V1998(01)
- (2) Anon; PATENT ABSTRACTS OF JAPAN 2000, V2000(02)
- (3) Bailey, T; WO 0010573 A 2000 HCAPLUS
- (4) Bayer Ag; WO 0226706 A 2002 HCAPLUS
- (5) Biediger, R; WO 9853790 A 1998 HCAPLUS
- (6) Carter, P; PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE UNITED
STATES OF AMERICA 2001, V98(21), P11879 HCAPLUS
- (7) Krajcik, R; WO 0162237 A 2001 HCAPLUS
- (8) Magnus, M; WO 02074752 A 2002 HCAPLUS
- (9) Ono Pharmaceut Co Ltd; JP 09249669 A 1997 HCAPLUS
- (10) Ono Pharmaceut Co Ltd; JP 11302280 A 1999 HCAPLUS
- (11) Wella Ag; DE 4027038 A 1992 HCAPLUS

IT 2295-31-0, 2,4-**Thiazolidinedione**

RL: RCT (Reactant); RACT (Reactant or reagent)
(use of heterocyclic compound or one of its salts to stimulate or induce
growth of hair and/or to slow down its **loss**)

RN 2295-31-0 HCAPLUS
 CN 2,4-Thiazolidinedione (8CI, 9CI) (CA INDEX NAME)



L97 ANSWER 2 OF 6 HCAPLUS COPYRIGHT 2004 ACS on STN
 AN 2002:978338 HCAPLUS
 DN 138:44664
 ED Entered STN: 29 Dec 2002
 TI Cosmetic compositions having retarding action on the regrowth of
 superfluous hair
 IN Di Pierro, Francesco
 PA Italy
 SO U.S. Pat. Appl. Publ., 6 pp., Cont.-in-part of U.S. Ser. No. 781,301,
 abandoned.
 CODEN: USXXCO

DT Patent
 LA English
 IC ICM A61K006-00
 ICS A61K007-00; A61K035-78
 NCL 424401000; 424725000
 CC 63-4 (Pharmaceuticals)
 FAN.CNT 2

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 2002197290	A1	20021226	US 2002-152805	20020523
	US 2001033849	A1	20011025	US 2001-781301	20010213
PRAI	IT 2000-MI628	A	20000324		
	US 2001-781301	B2	20010213		

AB The present invention relates to cosmetic compns. having retarding action
 on the regrowth of superfluous hair, more particularly to cosmetic compns.
 containing lipophilic exts. of Serenoa (Serenoa repens) enriched in fatty
 acids and with a reduced content of sterols. Preparation of Serenoa extract
 and

cosmetic preps. containing this extract is disclosed.

ST cosmetic **hair growth** inhibitor Serenoa ext

IT Anti-inflammatory agents

Cosmetics

Ginkgo biloba

Horse chestnut (Aesculus)

Horse chestnut (Aesculus hippocastanum)

Licorice (Glycyrrhiza glabra)

Ruscus aculeatus

Terminalia sericea

Vaccinium myrtillus

Zanthoxylum bungei

(cosmetic compns. having retarding action on regrowth of superfluous
 hair)

IT Anthocyanins

Fatty acids, biological studies

Flavones

Glycosides


Phospholipids, biological studies

Saponins

Terpenes, biological studies

Triterpenes

- RL: COS (Cosmetic use); BIOL (Biological study); USES (Uses)
(cosmetic compns. having retarding action on regrowth of superfluous hair)
- IT Cosmetics
(creams, depilatory; cosmetic compns. having retarding action on regrowth of superfluous hair)
- IT Cosmetics
(depilatories; cosmetic compns. having retarding action on regrowth of superfluous hair)
- IT Emulsions
Solutions
(depilatory; cosmetic compns. having retarding action on regrowth of superfluous hair)
- IT Thiols (organic), biological studies
RL: COS (Cosmetic use); BIOL (Biological study); USES (Uses)
(derivative; cosmetic compns. having retarding action on regrowth of superfluous hair)
- IT Serenoa repens
(exts.; cosmetic compns. having retarding action on regrowth of superfluous hair)
- IT Cosmetics
(gels, depilatory; cosmetic compns. having retarding action on regrowth of superfluous hair)
- IT **Hair preparations**
(**growth inhibitors**; cosmetic compns. having retarding action on regrowth of superfluous hair)
- IT Pruritus
(inhibitors; cosmetic compns. having retarding action on regrowth of superfluous hair)
- IT Cosmetics
(lotions, depilatory; cosmetic compns. having retarding action on regrowth of superfluous hair)
- IT Sterols
RL: COS (Cosmetic use); BIOL (Biological study); USES (Uses)
(phyto-; cosmetic compns. having retarding action on regrowth of superfluous hair)
- IT Phenols, biological studies
RL: COS (Cosmetic use); BIOL (Biological study); USES (Uses)
(polyphenols, nonpolymeric; cosmetic compns. having retarding action on regrowth of superfluous hair)
- IT Grape
(seeds; cosmetic compns. having retarding action on regrowth of superfluous hair)
- IT **56-03-1**, Biguanide 57-13-6, Urea, biological studies 60-23-1
62-56-6, Thiourea, biological studies 68-11-1, Thioglycolic acid, biological studies 79-42-5, Thiolactic acid 79-42-5D, Thiolactic acid, Alkaline-earth metal salts 126-97-6, Ethanolamine thioglycolate 507-09-5, Thioacetic acid, biological studies 563-83-7 7727-43-7, Barium sulfate
RL: COS (Cosmetic use); BIOL (Biological study); USES (Uses)
(cosmetic compns. having retarding action on regrowth of superfluous hair)
- IT **56-03-1**, Biguanide
RL: COS (Cosmetic use); BIOL (Biological study); USES (Uses)
(cosmetic compns. having retarding action on regrowth of superfluous hair)
- RN 56-03-1 HCAPLUS
- CN Imidodicarbonimidic diamide (9CI) (CA INDEX NAME)



L97 ANSWER 3 OF 6 HCAPLUS COPYRIGHT 2004 ACS on STN
 AN 2001:730517 HCAPLUS
 DN 135:277721
 ED Entered STN: 07 Oct 2001
 TI Cosmetic compositions containing antiandrogenic sterols with retarding
 action on the regrowth of superfluous hair
 IN Di Pierro, Francesco
 PA Indena S.P.A., Italy
 SO PCT Int. Appl., 17 pp.
 CODEN: PIXXD2
 DT Patent
 LA English
 IC ICM A61K007-06
 ICS A61K007-155
 CC 62-3 (Essential Oils and Cosmetics)
 FAN.CNT 2

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2001072266	A1	20011004	WO 2001-EP1522	20010212
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
	EP 1265586	A1	20021218	EP 2001-909738	20010212
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR				
	JP 2003528123	T2	20030924	JP 2001-570228	20010212
	NO 2002004519	A	20021125	NO 2002-4519	20020920
PRAI	IT 2000-MI628	A	20000324		
	WO 2001-EP1522	W	20010212		
AB	The present invention relates to cosmetic compns. having retarding action on the regrowth of superfluous hair, more particularly to cosmetic compns. containing fatty acids and antiandrogenic sterols from serenoa (<i>Serenoa</i> <i>repens</i>) and/or from <i>Cucurbita</i> seeds (<i>Cucurbita pepo</i>). A hair gel contained <i>Serenoa repens</i> lipophilic extract 2.00, ruscogenins 0.30, 20% <i>zanthoxylum bungenanum</i> extract 0.50, ethanol 20.00, Softigen-767 15.00, propylene glycol 10.00, Oleth-20 5.00, dimethicone copolyol 2.50, carbomer 2.00, triethanolamine 1.00, zinc ricinoleate 0.20, menthol 0.50, preservatives q.s., antioxidants q.s., and water q.s. 100.00 g.				
ST	cosmetic hair growth inhibitor antiandrogenic sterol				
IT	<i>Vaccinium myrtillus</i> (anthocyanosides; cosmetic compns. containing antiandrogenic sterols with retarding action on regrowth of superfluous hair)				
IT	Androgens RL: BOC (Biological occurrence); BSU (Biological study, unclassified); BUU (Biological use, unclassified); BIOL (Biological study); OCCU (Occurrence); USES (Uses) (antiandrogens; cosmetic compns. containing antiandrogenic sterols with retarding action on regrowth of superfluous hair)				
IT	Analgesics Anti-inflammatory agents <i>Licorice</i> (<i>Glycyrrhiza</i>) <i>Serenoa</i> <i>Serenoa repens</i> <i>Squash</i> (<i>Cucurbita pepo</i>) <i>Terminalia sericea</i>				

- (cosmetic compns. containing antiandrogenic sterols with retarding action on regrowth of superfluous hair)
- IT Anthocyanins
Phospholipids, biological studies
Saponins
Sterols
Triterpenes
RL: BOC (Biological occurrence); BSU (Biological study, unclassified); BUU (Biological use, unclassified); BIOL (Biological study); OCCU (Occurrence); USES (Uses)
(cosmetic compns. containing antiandrogenic sterols with retarding action on regrowth of superfluous hair)
- IT Alkali metal salts
Alkaline earth salts
Thiols (organic), biological studies
RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)
(cosmetic compns. containing antiandrogenic sterols with retarding action on regrowth of superfluous hair)
- IT Cosmetics
(creams; cosmetic compns. containing antiandrogenic sterols with retarding action on regrowth of superfluous hair)
- IT Cosmetics
(depilatories; cosmetic compns. containing antiandrogenic sterols with retarding action on regrowth of superfluous hair)
- IT Cosmetics
(emulsions; cosmetic compns. containing antiandrogenic sterols with retarding action on regrowth of superfluous hair)
- IT Glycosides
RL: BOC (Biological occurrence); BSU (Biological study, unclassified); BUU (Biological use, unclassified); BIOL (Biological study); OCCU (Occurrence); USES (Uses)
(flavonoid, oxo; cosmetic compns. containing antiandrogenic sterols with retarding action on regrowth of superfluous hair)
- IT Ginkgo biloba
(flavonoids from; cosmetic compns. containing antiandrogenic sterols with retarding action on regrowth of superfluous hair)
- IT Cosmetics
(foams; cosmetic compns. containing antiandrogenic sterols with retarding action on regrowth of superfluous hair)
- IT Cosmetics
(gels; cosmetic compns. containing antiandrogenic sterols with retarding action on regrowth of superfluous hair)
- IT **Hair preparations**
(**growth inhibitors**; cosmetic compns. containing antiandrogenic sterols with retarding action on regrowth of superfluous hair)
- IT Edema
(inhibitors; cosmetic compns. containing antiandrogenic sterols with retarding action on regrowth of superfluous hair)
- IT Zanthoxylum bungei
(isobutylamides from; cosmetic compns. containing antiandrogenic sterols with retarding action on regrowth of superfluous hair)
- IT Cosmetics
(lotions; cosmetic compns. containing antiandrogenic sterols with retarding action on regrowth of superfluous hair)
- IT Phenols, biological studies
RL: BOC (Biological occurrence); BSU (Biological study, unclassified); BUU (Biological use, unclassified); BIOL (Biological study); OCCU (Occurrence); USES (Uses)
(polyphenols, nonpolymeric; cosmetic compns. containing antiandrogenic sterols with retarding action on regrowth of superfluous hair)
- IT Grape

(polyphenols; cosmetic compns. containing antiandrogenic sterols with retarding action on regrowth of superfluous hair)

IT Horse chestnut (*Aesculus hippocastanum*)
Ruscus aculeatus
 (saponins; cosmetic compns. containing antiandrogenic sterols with retarding action on regrowth of superfluous hair)

IT Cucurbita
 (seeds; cosmetic compns. containing antiandrogenic sterols with retarding action on regrowth of superfluous hair)

IT 471-53-4, glycyrrhetic acid 472-11-7, ruscogenin 563-83-7
 RL: BOC (Biological occurrence); BSU (Biological study, unclassified); BUU (Biological use, unclassified); BIOL (Biological study); OCCU (Occurrence); USES (Uses)
 (cosmetic compns. containing antiandrogenic sterols with retarding action on regrowth of superfluous hair)

IT 56-03-1D, Biguanide, derivs. 60-23-1, 2-Aminoethanethiol
 62-56-6, Thiourea, biological studies 68-11-1, Thioglycolic acid, biological studies 79-42-5, Thiolactic acid 126-97-6, Ethanolamine thioglycolate 507-09-5, Thioacetic acid, biological studies 7727-43-7, Bariumsulfate 9002-13-5, Urease 30232-12-3, Mercaptopropionic acid
 RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)
 (cosmetic compns. containing antiandrogenic sterols with retarding action on regrowth of superfluous hair)


RE.CNT 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD

RE
 (1) Barr, E; WO 9833472 A 1998 HCAPLUS
 (2) Greentech S A; FR 2791255 A 2000

IT 56-03-1D, Biguanide, derivs.
 RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)
 (cosmetic compns. containing antiandrogenic sterols with retarding action on regrowth of superfluous hair)

RN 56-03-1 HCAPLUS

CN Imidodicarbonimidic diamide (9CI) (CA INDEX NAME)



L97 ANSWER 4 OF 6 HCAPLUS COPYRIGHT 2004 ACS on STN

AN 2001:635880 HCAPLUS

DN 135:200473

ED Entered STN: 31 Aug 2001

TI Methods and compositions based on **insulin-sensitivity increasing substances** for the treatment of **alopecia** and other disorders of the **pilosebaceous apparatus**

IN **Krajcik, Rozlyn A.; Orentreich, Norman**

PA Orentreich Foundation for the Advancement of Science, Inc., USA

SO PCT Int. Appl., 22 pp.
 CODEN: PIXXD2

DT Patent

LA English

IC ICM A61K031-00

CC 63-6 (Pharmaceuticals)
 Section cross-reference(s): 1, 2, 62

FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	-----	-----	-----	-----


PI WO 2001062237 A2 20010830 WO 2001-US5653 20010223 <--
 WO 2001062237 A3 20020613
 W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM
 RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG
 EP 1267850 A2 20030102 EP 2001-914437 20010223 <--
 R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR
 US 2002143039 A1 20021003 US 2002-73607 20020211 <--
 PRAI US 2000-184398P P 20000223 <--
 WO 2001-US5653 W 20010223 <--
 AB **Insulin sensitivity increasing substances (ISIS)**, including but not limited to **D-chiro-inositol, thiazolidinedione** and derivs., and biguanides, are useful in the treatment of **hair loss** and other disorders of the **pilosebaceous** apparatus (hirsutism, acne, etc.) associated with conditions of excess **insulin** and/or **insulin** resistance. The treatment comprises administering to a mammal, such as a human, at least one **ISIS** either alone or in combination with at least one agent, such as an androgen receptor blocker (ARB) and/or a steroid enzyme inhibitor or inducer (STI). Addnl., an activity enhancing agent may be included for topical administration. For example, the onset of age-dependent **hair loss** in female ob/ob (obese) mice was delayed by oral **metformin-HCl** treatment using a dose of 240 mg/kg. Clear differences were seen between the incidence of **hair loss** in control vs. **metformin HCl**-treated animals in animals that were older than 300 days. The incidence of **hair loss** in **metformin HCl**-treated animals at 370 days of age was 30% compared to 60% incidence of **hair loss** in non-treated animals. In animals that were 300 days of age, about 20% of the **metformin HCl**-treated animals exhibited **hair loss** in contrast to the control animals, which showed about a 40% incidence of **hair loss**. Addnl., it was noted in the study that obese mice were prone to a spontaneous skin condition which may resemble human acanthosis nigricans or migratory ichthyosis. Although this condition was not fully characterized, the **metformin HCl**-treated animal group exhibited markedly less incidence of this skin condition relative to the control animals, the majority of which were affected by the skin condition. In addition, transient changes in **hair loss** patterns were occasionally noted in some of the animals during the course of the study. For example, an animal which presented with very moderate **hair loss** (i.e., only possible thinning of **hair** coat) for a period of 2-3 wk might later exhibit no **hair loss** and sustain that grade for an extended period of time.
 ST biguanide inositol **thiazolidinedione** insulin sensitivity **alopecia**; oral biguanide inositol **thiazolidinedione** **alopecia**; topical biguanide inositol **thiazolidinedione** **alopecia**; **hair growth promoter** biguanide inositol **thiazolidinedione**
 IT Skin, disease
 (acanthosis nigricans; compns. containing insulin-sensitivity increasing compds. for treatment of **alopecia** and other disorders of **pilosebaceous** apparatus)
 IT Androgen receptors
 RL: BSU (Biological study, unclassified); BIOL (Biological study)

(blockers; compns. containing insulin-sensitivity increasing compds. for treatment of **alopecia** and other disorders of **pilosebaceous** apparatus)


- IT Acne
 - Alopecia**
 - Anti-inflammatory agents
 - Antidiabetic agents
 - Hirsutism**
 - Permeation enhancers
 - Surfactants
 - Vasodilators
 - (compns. containing insulin-sensitivity increasing compds. for treatment of **alopecia** and other disorders of **pilosebaceous** apparatus)
- IT Drug delivery systems
 - (gels; compns. containing insulin-sensitivity increasing compds. for treatment of **alopecia** and other disorders of **pilosebaceous** apparatus)
- IT **Hair preparations**
 - (**growth stimulants**; compns. containing insulin-sensitivity increasing compds. for treatment of **alopecia** and other disorders of **pilosebaceous** apparatus)
- IT Skin, disease
 - (ichthyosis, migratory; compns. containing insulin-sensitivity increasing compds. for treatment of **alopecia** and other disorders of **pilosebaceous** apparatus)
- IT Drug delivery systems
 - (ointments, creams; compns. containing insulin-sensitivity increasing compds. for treatment of **alopecia** and other disorders of **pilosebaceous** apparatus)
- IT Drug delivery systems
 - (oral; compns. containing insulin-sensitivity increasing compds. for treatment of **alopecia** and other disorders of **pilosebaceous** apparatus)
- IT Enzymes, biological studies
 - RL: BSU (Biological study, unclassified); BIOL (Biological study)
 - (steroidogenic, inhibitors or inducers; compns. containing insulin-sensitivity increasing compds. for treatment of **alopecia** and other disorders of **pilosebaceous** apparatus)
- IT Drug delivery systems
 - (tinctures; compns. containing insulin-sensitivity increasing compds. for treatment of **alopecia** and other disorders of **pilosebaceous** apparatus)
- IT Drug delivery systems
 - (topical; compns. containing insulin-sensitivity increasing compds. for treatment of **alopecia** and other disorders of **pilosebaceous** apparatus)
- IT 67-68-5, Dimethyl sulfoxide, biological studies 872-50-4,
 N-Methylpyrrolidone, biological studies 3079-28-5, Decylmethyl sulfoxide
 63839-46-3, Myristylamine oxide
 RL: MOA (Modifier or additive use); THU (Therapeutic use); BIOL
 (Biological study); USES (Uses)
 (compns. containing insulin-sensitivity increasing compds. for treatment of **alopecia** and other disorders of **pilosebaceous** apparatus)
- IT 52-01-7, Spironolactone 56-03-1D, Biguanide,
 derivs. 57-83-0, Progesterone, biological studies
 102-02-3, Phenyl biguanide 427-51-0, Cyproterone
 acetate 643-12-9, D-Chiro-
 Inositol 976-71-6, Canrenone 1115-70-4
 2295-31-0D, Thiazolidinedione, derivs.
 13311-84-7, Flutamide 34461-22-8,
 Metformin pamoate 51481-61-9, Cimetidine
 63612-50-0, Nilutamide 65277-42-1,
 Ketoconazole 73671-86-0, 4-MA

- 74772-77-3, Ciglitazone 90357-06-5,
Bicalutamide 97322-87-7, Troglitazone
109229-58-5, Englitazone 111025-46-8,
Pioglitazone 122320-73-4, Rosiglitazone
154992-24-2, RU-58841
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(compns. containing insulin-sensitivity increasing compds. for treatment of
alopecia and other disorders of **pilosebaceous** apparatus)
- IT 9015-81-0
RL: BSU (Biological study, unclassified); BIOL (Biological study)
(inducers; compns. containing insulin-sensitivity increasing compds. for
treatment of **alopecia** and other disorders of
pilosebaceous apparatus)
- IT 9028-56-2, 3 α -Hydroxy-
steroid dehydrogenase 9081-34-9, 5
 α -Reductase
RL: BSU (Biological study, unclassified); BIOL (Biological study)
(inhibitors; compns. containing insulin-sensitivity increasing compds. for
treatment of **alopecia** and other disorders of
pilosebaceous apparatus)
- IT 50-99-7, D-Glucose, biological studies
RL: BSU (Biological study, unclassified); BIOL (Biological study)
(regulating agents; compns. containing insulin-sensitivity increasing
compds. for treatment of **alopecia** and other disorders of
pilosebaceous apparatus)
- IT 9004-10-8, Insulin, biological studies
RL: BSU (Biological study, unclassified); BIOL (Biological study)
(sensitivity; compns. containing insulin-sensitivity increasing compds. for
treatment of **alopecia** and other disorders of
pilosebaceous apparatus)
- IT 52-01-7, Spironolactone 56-03-1D, Biguanide,
derivs. 57-83-0, Progesterone, biological studies
102-02-3, Phenyl biguanide 427-51-0, Cyproterone
acetate 643-12-9, D-Chiro-
Inositol 976-71-6, Canrenone 1115-70-4
2295-31-0D, Thiazolidinedione, derivs.
13311-84-7, Flutamide 34461-22-8,
Metformin pamoate 51481-61-9, Cimetidine
63612-50-0, Nilutamide 65277-42-1,
Ketoconazole 73671-86-0, 4-MA
74772-77-3, Ciglitazone 90357-06-5,
Bicalutamide 97322-87-7, Troglitazone
109229-58-5, Englitazone 111025-46-8,
Pioglitazone 122320-73-4, Rosiglitazone
154992-24-2, RU-58841
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(compns. containing insulin-sensitivity increasing compds. for treatment of
alopecia and other disorders of **pilosebaceous** apparatus)
- RN 52-01-7 HCAPLUS
- CN Pregn-4-ene-21-carboxylic acid, 7-(acetylthio)-17-hydroxy-3-oxo-,
 γ -lactone, (7 α ,17 α)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 56-03-1 HCAPLUS
 CN Imidodicarbonimidic diamide (9CI) (CA INDEX NAME)




RN 57-83-0 HCAPLUS
 CN Pregn-4-ene-3,20-dione (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 102-02-3 HCAPLUS
 CN Imidodicarbonimidic diamide, N-phenyl- (9CI) (CA INDEX NAME)




RN 427-51-0 HCAPLUS
 CN 3'H-Cyclopropa[1,2]pregna-1,4,6-triene-3,20-dione, 17-(acetyloxy)-6-chloro-1,2-dihydro-, (1 β ,2 β)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 643-12-9 HCAPLUS
 CN D-chiro-Inositol (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 976-71-6 HCAPLUS
 CN Pregna-4,6-diene-21-carboxylic acid, 17-hydroxy-3-oxo-, γ -lactone, (17 α)- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (+).




RN 1115-70-4 HCAPLUS
 CN Imidodicarbonimidic diamide, N,N-dimethyl-, monohydrochloride (9CI) (CA INDEX NAME)




● HCl

RN 2295-31-0 HCAPLUS
 CN 2,4-Thiazolidinedione (8CI, 9CI) (CA INDEX NAME)




RN 13311-84-7 HCAPLUS
 CN Propanamide, 2-methyl-N-[4-nitro-3-(trifluoromethyl)phenyl]- (9CI) (CA INDEX NAME)



RN 34461-22-8 HCAPLUS
 CN 2-Naphthalenecarboxylic acid, 4,4'-methylenebis[3-hydroxy-, compd. with N,N-dimethylimidodicarbonimidic diamide (1:2) (9CI) (CA INDEX NAME)


CM 1

CRN 657-24-9
 CMF C4 H11 N5




CM 2

CRN 130-85-8
 CMF C23 H16 O6




RN 51481-61-9 HCAPLUS

CN Guanidine, N-cyano-N'-methyl-N''-[2-[[[5-methyl-1H-imidazol-4-yl)methyl]thio]ethyl]- (9CI) (CA INDEX NAME)



RN 63612-50-0 HCAPLUS


CN 2,4-Imidazolidinedione, 5,5-dimethyl-3-[4-nitro-3-(trifluoromethyl)phenyl]- (9CI) (CA INDEX NAME)



RN 65277-42-1 HCAPLUS

CN Piperazine, 1-acetyl-4-[4-[[[(2R,4S)-2-(2,4-dichlorophenyl)-2-(1H-imidazol-1-ylmethyl)-1,3-dioxolan-4-yl]methoxy]phenyl]-, rel- (9CI) (CA INDEX NAME)


Relative stereochemistry.



RN 73671-86-0 HCAPLUS


CN 1H-Indeno[5,4-f]quinoline-7-carboxamide, N,N-diethylhexadecahydro-1,4a,6a-trimethyl-2-oxo-, (4aR,4bS,6aS,7S,9aS,9bS,11aR) - (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 74772-77-3 HCAPLUS

CN 2,4-Thiazolidinedione, 5-[[4-[(1-methylcyclohexyl)methoxy]phenyl]methyl] - (9CI) (CA INDEX NAME)




RN 90357-06-5 HCAPLUS

CN Propanamide, N-[4-cyano-3-(trifluoromethyl)phenyl]-3-[(4-fluorophenyl)sulfonyl]-2-hydroxy-2-methyl- (9CI) (CA INDEX NAME)




RN 97322-87-7 HCAPLUS

CN 2,4-Thiazolidinedione, 5-[[4-[(3,4-dihydro-6-hydroxy-2,5,7,8-tetramethyl-2H-1-benzopyran-2-yl)methoxy]phenyl]methyl]- (9CI) (CA INDEX NAME)



RN 109229-58-5 HCAPLUS


CN 2,4-Thiazolidinedione, 5-[[3,4-dihydro-2-(phenylmethyl)-2H-1-benzopyran-6-yl]methyl]- (9CI) (CA INDEX NAME)



RN 111025-46-8 HCAPLUS

CN 2,4-Thiazolidinedione, 5-[[4-[2-(5-ethyl-2-pyridinyl)ethoxy]phenyl]methyl]-
(9CI) (CA INDEX NAME)

PAGE 1-A




PAGE 2-A

Et


RN 122320-73-4 HCAPLUS

CN 2,4-Thiazolidinedione, 5-[[4-[2-(methyl-2-pyridinylamino)ethoxy]phenyl]met
hyl]- (9CI) (CA INDEX NAME)


PAGE 1-A



PAGE 2-A



RN 154992-24-2 HCAPLUS
 CN Benzonitrile, 4-[3-(4-hydroxybutyl)-4,4-dimethyl-2,5-dioxo-1-imidazolidinyl]-2-(trifluoromethyl)- (9CI) (CA INDEX NAME)



IT 9015-81-0
 RL: BSU (Biological study, unclassified); BIOL (Biological study)
 (inducers; compns. containing insulin-sensitivity increasing compds. for
 treatment of **alopecia** and other disorders of
pilosebaceous apparatus)

RN 9015-81-0 HCAPLUS
 CN Dehydrogenase, 3(17) β -hydroxy steroid (9CI) (CA INDEX NAME)

*** STRUCTURE DIAGRAM IS NOT AVAILABLE ***

IT 9028-56-2, 3 α -Hydroxy-
 steroid dehydrogenase 9081-34-9, 5
 α -Reductase

RL: BSU (Biological study, unclassified); BIOL (Biological study)
 (inhibitors; compns. containing insulin-sensitivity increasing compds. for
 treatment of alopecia and other disorders of
 pilosebaceous apparatus)

RN 9028-56-2 HCAPLUS
 CN Dehydrogenase, 3 α -hydroxy steroid (9CI) (CA INDEX NAME)

*** STRUCTURE DIAGRAM IS NOT AVAILABLE ***

RN 9081-34-9 HCAPLUS
 CN Reductase, steroid 5 α - (9CI) (CA INDEX NAME)

*** STRUCTURE DIAGRAM IS NOT AVAILABLE ***

L97 ANSWER 5 OF 6 HCAPLUS COPYRIGHT 2004 ACS on STN
 AN 2001:247151 HCAPLUS
 DN 134:271051
 ED Entered STN: 06 Apr 2001
 TI Cosmetic composition based on organic silicon compounds comprising at
 least a function with cosmetic effect
 IN Samain, Henri; Rollat, Isabelle; Jeanne Rose, Valerie; Sanchez, Clement
 PA L'oreal, Fr.
 SO PCT Int. Appl., 17 pp.
 CODEN: PIXXD2
 DT Patent
 LA French
 IC ICM A61K007-48
 ICS A61K007-06
 CC 62-3 (Essential Oils and Cosmetics)
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2001022932	A1	20010405	WO 1999-FR2291	19990927
	W:				
	AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU,				
	CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL,				
	IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD,				
	MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK,				
	SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ,				
	BY, KG, KZ, MD, RU, TJ, TM				
	RW:				
	GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE,				
	DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF,				
	CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
	AU 9958680	A1	20010430	AU 1999-58680	19990927
	BR 9917501	A	20020521	BR 1999-17501	19990927
	EP 1216023	A1	20020626	EP 1999-946239	19990927
	R:				
	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,				
	IE, SI, LT, LV, FI, RO, MK, CY, AL				
	JP 2003510262	T2	20030318	JP 2001-526144	19990927
PRAI	WO 1999-FR2291	A	19990927		

AB The invention concerns a composition comprising, in a cosmetically acceptable medium, at least 0.02 weight % relative to the composition total weight, one or several hardly or non-polymerized water soluble organic silicon compds., selected

among silanes comprising a silicon atom and siloxanes comprising two silicon atoms, said organic silicon compds. comprising per mol. two hydroxyl groups or capable of being hydrolyzed and two groups not capable of being hydrolyzed, at least one of said groups being a group with cosmetic effect

and at least one of the remaining functional groups not capable of being hydrolyzed being a group with a solubilizing function. The invention is applicable to hair care compns. A hair dye preparation contained aminopropyl-N-(4,2-dinitrophenyl)aminopropyl diethoxysilane 2, and a hydroalcoholic mixture (50:50) 98 g. The preparation was applied on a 70%

white

hair, then washed and dried to obtain an orange color.

ST cosmetic hair dye org silicon compd

IT Anthraquinone dyes

Azo dyes

Fungicides

Hair preparations

Reducing agents

Sunscreens

(cosmetic composition based on organic silicon compds. comprising at least function with cosmetic effect)

IT Polysiloxanes, biological studies

RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)

(cosmetic composition based on organic silicon compds. comprising at least function with cosmetic effect)

IT Antibiotics

(macrolide; cosmetic composition based on organic silicon compds.

comprising at

least function with cosmetic effect)

IT 56-03-1, Biguanide 268724-97-6

RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)

(cosmetic composition based on organic silicon compds. comprising at least function with cosmetic effect)

RE.CNT 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD

RE

(1) Meyer, T; US 5750092 A 1998 HCAPLUS

(2) Shiseido Co Ltd; EP 0655453 A 1995 HCAPLUS


IT 56-03-1, Biguanide

RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)

(cosmetic composition based on organic silicon compds. comprising at least function with cosmetic effect)

RN 56-03-1 HCAPLUS

CN Imidodicarbonimidic diamide (9CI) (CA INDEX NAME)



L97 ANSWER 6 OF 6 HCAPLUS COPYRIGHT 2004 ACS on STN

AN 1998:65452 HCAPLUS

DN 128:145146

ED Entered STN: 04 Feb 1998

TI Hair treatment compositions

IN Kashino, Takayoshi; Nagai, Minoru; Ono, Toshinari; Tabata, Yoshiko; Hirano, Aya

PA Kao Corp., Japan

SO Jpn. Kokai Tokkyo Koho, 13 pp.

CODEN: JKXXAF

DT Patent

LA Japanese


IC ICM A61K007-09

ICS A61K007-00

CC 62-3 (Essential Oils and Cosmetics)

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	JP 10017442	A2	19980120	JP 1996-174774	19960704
PRAI	JP 1996-174774		19960704		
OS	MARPAT 128:145146				
AB	Hair treatment compns. showing no damage to hair contain: (A) keratin reduction substances such as thioglycolic acid and N-acetyl-L-cysteine and (B) guanidine derivs. such as butylguanidine and hexylguanidine, in addition to other ingredients.				
ST	hair prepn guanidine deriv; keratin redn substance hair prepn; thioglycolic acid guanidine deriv hair prepn				
IT	Hair preparations (hair treatment compns.)				
IT	Keratins RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses) (reduction substances; hair treatment compns.)				
IT	68-11-1, Thioglycolic acid, biological studies 80-70-6, 1,1,3,3-Tetramethylguanidine 102-02-3, 1-Phenylbiguanide 113-00-8D, Guanidine, derivs. 462-69-1, Butylguanidine 543-18-0, 2-Guanidinoethanesulfonic acid 616-91-1, N-Acetyl-L-cysteine 692-13-7, 1-Butylbiguanide 1119-69-3 1866-59-7, Hexylguanidine 2002-16-6, Phenylguanidine 3324-71-8, 1,3-Dimethylguanidine 14948-83-5, Cyclohexylguanidine 20600-59-3 48138-07-4 67337-40-0 89282-88-2 100056-66-4 139419-37-7 176370-21-1 202396-97-2 RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses) (hair treatment compns.)				
IT	102-02-3, 1-Phenylbiguanide RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses) (hair treatment compns.)				
RN	102-02-3 HCAPLUS				
CN	Imidodicarbonimidic diamide, N-phenyl- (9CI) (CA INDEX NAME)				



=> d 182 bib abs hitrn tot

L82 ANSWER 1 OF 11 HCAPLUS COPYRIGHT 2004 ACS on STN

AN 1994:455727 HCAPLUS

DN 121:55727

TI Antibodies to **hair follicles** in **alopecia areata**

AU Tobin, Desmond J.; Orentreich, Norman; Fenton, David A.; Bystryn, Jean-Claude

CS Dep. Dermatol., New York Univ. Med. Cent., New York, NY, 10016, USA

SO Journal of Investigative Dermatology (1994), 102(5), 721-4

CODEN: JIDEAE; ISSN: 0022-202X

DT Journal

LA English

AB Although **alopecia areata** is suspected to be an autoimmune disease, no direct evidence of an altered immune response to components of the **hair follicle** has been reported. The authors studied whether antibodies to normal human anagen **scalp**

hair follicles are present in individuals with **alopecia areata**. Thirty-nine **alopecia areata** sera and 27 control sera were tested by Western immunoblotting for antibodies to 6 M urea-extractable proteins of normal anagen **scalp hair follicles**. At serum diluted 1:80, all **alopecia areata** subjects (100%), but only 44% of control individuals, had antibodies directed to one or more antigens of .apprx.57, 52, 50, 47, or 44 kDa. The incidence of antibodies to individual **hair follicle** antigens in **alopecia areata** was up to seven times more frequent than in control sera and their level up to 13 times greater and was significant for all five antigens. Tissue specificity anal. indicated that these antigens were selectively expressed in **hair follicles**. These findings indicate that individuals with **alopecia areata** have abnormal antibodies directed to **hair follicle** antigens, and support the hypothesis that **alopecia areata** is an autoimmune disease.

L82 ANSWER 2 OF 11 HCAPLUS COPYRIGHT 2004 ACS on STN

AN 1989:69565 HCAPLUS

DN 110:69565

TI Local stimulation of sebaceous gland activity by the topical administration of dehydroepiandrosterone

AU Orentreich, Norman; Matias, Jonathan R.

CS Biomed. Res. Stn., Orentreich Found. Adv. Sci., Inc., Cold Spring, NY, 10516, USA

SO Journal of the Society of Cosmetic Chemists (1988), 39(5), 291-303
CODEN: JSCCA5; ISSN: 0037-9832

DT Journal

LA English

AB Topical preps. of dehydroepiandrosterone (DHEA), were tested exptl. in laboratory animals and clin. in women. When applied topically at a concentration of

1%, DHEA stimulated **growth** of the ventral ear skin sebaceous glands of female Syrian hamsters by .apprx.80% after 2 wk of daily application without any evidence of systemic effects. Dose-response studies demonstrated that topical DHEA was effective at concns. as low as 0.1%. Systemic side effects, as measured by gland size of the contralateral ear, were absent even with the concns. <5% or with applications of 1% DHEA \leq 3 times per day. Topical DHEA did not induce hirsutism since **hair growth**-promoting effects were not observed in the androgen-dependent **hair** of the long-haired Syrian hamster. Sebum excretion rate was also measured in women over 1.5 yr on alternating treatments of placebo and DHEA creams. Topical DHEA doubled sebum production in women at a min. ED of 0.1%, without any evidence of untoward side effects. Thus, the topical application of DHEA is safe and effective in stimulating sebum production in women with reduced sebum output usually associated with menopause.

IT 57-83-0, Pregn-4-ene-3,20-dione, biological studies

RL: BIOL (Biological study)

(dehydroepiandrosterone-stimulated sebaceous gland growth inhibition by)

L82 ANSWER 3 OF 11 HCAPLUS COPYRIGHT 2004 ACS on STN

AN 1988:622776 HCAPLUS

DN 109:222776

TI Synergistic antiandrogenic effects of topical combinations of 5. **alpha.-reductase** and androgen receptor inhibitors in the hamster sebaceous glands

AU Matias, Jonathan R.; Malloy, Virginia L.; Orentreich, Norman

CS Biomed. Res. Stn., Orentreich Found. Adv. Sci., Inc., Cold Spring-on-Hudson, NY, 10516, USA

SO Journal of Investigative Dermatology (1988), 91(5), 429-33
CODEN: JIDEAE; ISSN: 0022-202X

DT Journal

LA English

AB The androgenic action of dihydrotestosterone (DHT) is antagonized by agents that compete with testosterone for the 5.alpha.-reductase enzyme and by agents that block the binding of DHT to its receptor. The topical synergistic effect of 5.alpha.-reductase (5 α RI) and androgen receptor inhibitors (ARI) was determined by measurement of the sebaceous gland size (SGS) of the ventral ear skin of the intact, sexually mature male Syrian hamsters. Progesterone, a 5 α RI, and spironolactone (SL), an ARI, produced a dose responsive decrease in SGS at topical concns. of 0.01-5.0%. At concns. of 1, 3, and 5%, progesterone and SL combinations produced neither an additive nor synergistic inhibition of SGS. At very low concns. of up to 0.10%, neither progesterone nor SL alone produced any effect on SGS. When combinations of these 2 steroids were applied at low concns., SGS decreased unilaterally to approx. 50%. This synergy occurred best at a P:SL ratio of 1:2. The lower effective concns. of progesterone may be explained by its greater percutaneous absorption. Synergy was also demonstrated at low concns. with other antiandrogens: cyproterone acetate, canrenone, hydroxyflutamide, and N,N-diethyl-4-methyl-3-oxo-4-aza-5- α -androstane-17 β -carboxamide. The use of antiandrogen combinations at low concns. is of value because of the decreased risk of systemic side effects while maintaining potent topical efficacy.

IT 427-51-0, Cyproterone acetate

RL: BIOL (Biological study)

(sebaceous gland size decrease by progesterone and)

IT 73671-86-0

RL: BIOL (Biological study)

(sebaceous gland size decrease by spironolactone and)

IT 52-01-7, Spironolactone 976-71-6,

Canrenone

RL: BIOL (Biological study)

(sebaceous gland size decrease by, progesterone synergism with)

IT 57-83-0, Progesterone, biological studies

RL: BIOL (Biological study)

(sebaceous gland size decrease by, spironolactone synergism with)

L82 ANSWER 4 OF 11 HCAPLUS COPYRIGHT 2004 ACS on STN

AN 1988:26977 HCAPLUS

DN 108:26977

TI Topical compositions containing androgen inhibitors for the treatment of sebaceous gland hypertrophy

IN Orentreich, Norman; Matias, Jonathan R.

PA USA

SO U.S., 4 pp. Cont. of U.S. Ser. No. 609,152, abandoned.

CODEN: USXXAM

DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 4684635	A	19870804	US 1986-846498	19860327
	CA 1243609	A1	19881025	CA 1985-481315	19850510
	US 5053403	A	19911001	US 1989-438979	19891120
PRAI	US 1984-609152		19840511		
	US 1986-846498		19860327		
	US 1987-79609		19870730		

AB A topical composition for the treatment of sebaceous gland hypertrophy and hirsutism comprises synergistically effective amts. of (1) an inhibitor of

the conversion of testosterone to dihydrotestosterone by the 5.
alpha.-reductase and (2) a blocking agent which blocks
the binding of dihydrotestosterone to receptor protein in cell cytoplasm.
Animal studies showed that the combination of 5.**alpha**
.-reductase inhibitor and androgen inhibitor produced a
synergistic effect. A topical solution contained **progesterone**
0.025, **spironolactone** 0.05% weight/volume and acetone to 100%
weight/volume

IT 52-01-7, **Spironolactone** 427-51-0,
Cyproterone acetate 976-71-6,
Canrenone 13311-84-7, **Flutamide**
63612-50-0

RL: BIOL (Biological study)
(sebaceous gland hypertrophy treatment with, in combination with
5 α -**reductase** inhibitors)

IT 57-83-0, **Progesterone**, biological studies

RL: BIOL (Biological study)
(sebaceous gland hypertrophy treatment with, in combination with
androgen receptor blocker)

L82 ANSWER 5 OF 11 HCAPLUS COPYRIGHT 2004 ACS on STN

AN 1985:400787 HCAPLUS

DN 103:787

TI A comparative study of the effects of percutaneously administered
5 α -**reductase** and androgen receptor
inhibitors on hamster sebaceous glands

AU Matias, Jonathan R.; Malloy, Virginia; **Orentreich, Norman**

CS Orentreich Found. Adv. Sci., Inc., New York, NY, 10021, USA

SO Annals of the New York Academy of Sciences (1984), 435(Colloq. Biol. Sci.,
1st, 1983), 454-6

CODEN: ANYAA9; ISSN: 0077-8923

DT Journal

LA English

AB The inhibition of the 5 α -reduction of testosterone by
progesterone (P) [57-83-0] and the inhibition of
dihydrotestosterone binding to the cytosol receptor by **cyproterone**
acetate (CA) [427-51-0] or **spironolactone**
(SL) [52-01-7] are important approaches for the treatment of
androgen-related cutaneous disorders. The effects of these steroids on
the size of the androgen-sensitive sebaceous glands of mature male Syrian
hamsters were studied. These compds. were dissolved in Me₂CO and applied
unilaterally on the ventral ear skin at doses ranging 50 μ g-2.5 mg/day
for 4 wk. A dose-related reduction of sebaceous gland size was shown for each
antiandrogen. At the highest dose, the magnitudes of the inhibitory
effect of P, CA, and SL were 58%, 40%, and 31%, resp. The antiandrogenic
activity of P was localized at the treated skin, whereas CA and SL caused
significant systemic side effects. Evidently, P is a more suitable
topical androgen antagonist than CA and SL because of its greater potency
and unilateral activity.

IT 52-01-7 427-51-0

RL: BAC (Biological activity or effector, except adverse); BSU (Biological
study, unclassified); BIOL (Biological study)
(sebaceous gland response to, androgen receptors inhibition in relation
to)

IT 57-83-0, biological studies

RL: BAC (Biological activity or effector, except adverse); BSU (Biological
study, unclassified); BIOL (Biological study)
(sebaceous gland response to, reductase inhibition in relation to)

L82 ANSWER 6 OF 11 HCAPLUS COPYRIGHT 2004 ACS on STN

AN 1984:604533 HCAPLUS


DN 101:204533

TI The local antiandrogenic effect of the intracutaneous injection of

progesterone in the flank organ of sexually mature male Syrian golden hamster

- AU **Orentreich, N.**; Matias, J. R.; Malloy, V.
 CS Med. Cent., New York Univ., New York, NY, 10016, USA
 SO Archives of Dermatological Research (1984), 276(6), 401-5
 CODEN: ADREDL; ISSN: 0340-3696
 DT Journal
 LA English
 AB The local antiandrogenic action of **progesterone** [57-83-0] was investigated using the androgen-responsive flank organs of adult, sexually mature male Syrian golden hamsters. The effects of unilateral intracutaneous injections of a micronized suspension of **progesterone** into the flank organs was analyzed by the measurement of the weight, area of surface pigmentation, and the cross-sectional area of the sebaceous glands. Weekly injections of 5 mg of **progesterone** for 3 wk produced approx. 50% reduction in all 3 parameters in comparison with the controls. The minimal ED of 1 mg/wk was determined by the injection of **progesterone** at doses of 0.1-5 mg. These effects were localized only to the treated flank organs, since the values for the contralateral side were not significantly different from control. The local action of **progesterone** was further demonstrated by the absence of effect on the weight of seminal vesicles and testes of the treated animals in comparison with the controls.
- IT 57-83-0, biological studies
 RL: BIOL (Biological study)
 (antiandrogenic activities of, in flank organ of male hamster)

- L82 ANSWER 7 OF 11 HCAPLUS COPYRIGHT 2004 ACS on STN
 AN 1984:563905 HCAPLUS
 DN 101:163905
 TI The lack of effect of 11 α - **hydroxyprogesterone** on the flank-organ and ear sebaceous glands of adult male Syrian golden hamsters
 AU Matias, J. R.; **Orentreich, N.**; Malloy, V.; De Feo, C. P., III; Matias, L.
 CS Anim. Sci. Lab., Orentreich Found. Adv. Sci., New York, NY, 10021, USA
 SO Archives of Dermatological Research (1984), 276(5), 346-8
 CODEN: ADREDL; ISSN: 0340-3696
 DT Journal
 LA English
 GI



- AB Topical application of 11 α - **hydroxyprogesterone** (I) [80-75-1] did not decrease the area of pigmentation of the flank organ or the size of the medial ear sebaceous glands, both of which were decreased by **progesterone** [57-83-0]. The lack of antiandrogenic activity of I raises questions as to the utility of this synthetic steroid for the treatment of androgen-mediated cutaneous diseases.
- IT 57-83-0, biological studies
 RL: BIOL (Biological study)
 (antiandrogenic activity of, **hydroxyprogesterone** in

comparison with)

L82 ANSWER 8 OF 11 HCAPLUS COPYRIGHT 2004 ACS on STN
AN 1984:96957 HCAPLUS
DN 100:96957
TI In vitro testosterone metabolism in the mouse preputial gland:
intercellular cooperation and changes with cell maturation
AU Brind, J. L.; Marinescu, D.; Gomez, E. C.; Wheatley, V. R.;
Orentreich, N.
CS Orentreich Found. Adv. Sci., New York, NY, 10021, USA
SO Journal of Endocrinology (1984), 100(3), 377-88
CODEN: JOENAK; ISSN: 0022-0795
DT Journal
LA English
AB In vitro ¹⁴C-labeled testosterone [58-22-0] metabolism was investigated in
isolated cells of adult male mouse preputial sebaceous glands. Labeled
steroids were extracted and chromatographed after a 2-h incubation and were
identified as 5 α -dihydrotestosterone [521-18-6], androstenedione
[63-05-8], 5 α -androstane-3,17-dione [846-46-8],
5 α -androstane-3 α ,17 β -diol [1852-53-5],
5 α -androstane-3 β ,17 β -diol [571-20-0], androsterone
[53-41-8], and 3-epiandrosterone [481-29-8]. In cells separated according to
state of maturity (lipid content) by isopycnic centrifugation in a
metrizamide gradient, maximal testosterone metabolism occurred in large nearly
mature cells. In this population mean hydroxy steroid 5.
alpha.-reductase [37325-56-7] and 17.
beta.-hydroxy steroid dehydrogenase
[9015-81-0] activities were 3.8 and 2.3 nmol/106 cells/2 h,
resp., >100-fold greater than in the densest population comprised of
undifferentiated and early differentiating cells. The profile of
testosterone metabolites depended on the proportion of the label
metabolized. The metabolite index (MI), i.e. the average number of enzymic
steps
undergone per mol. of metabolite, increased with increasing substrate
utilization. Metrizamide showed reversible, nonspecific inhibition of
testosterone metabolism and reduction of the MI. Thus, testosterone is
metabolized sequentially by different cells, with metrizamide inhibiting
cellular uptake and intercellular substrate transport. This suggested
that most of the metabolites would be found in the medium, rather than in
the cellular compartment. Further, in incubations run without cell
disaggregation, efficient substrate cycling among cells should result in a
high MI, independent of metrizamide concentration and substrate utilization.
These predictions were all confirmed, providing strong evidence that
testosterone metabolism is a cooperative effort among several cells in this
tissue.
IT 9015-81-0
RL: BOC (Biological occurrence); BSU (Biological study, unclassified);
BIOL (Biological study); OCCU (Occurrence)
(of Tyson's gland)

L82 ANSWER 9 OF 11 HCAPLUS COPYRIGHT 2004 ACS on STN
AN 1977:40706 HCAPLUS
DN 86:40706
TI Testosterone metabolism in human **scalp** and beard **hair**
follicles
AU Rizer, Ronald L.; **Orentreich, Norman**; Finch, Caleb E.
CS Orentreich Med. Group, New York, NY, USA
SO Hum. Hair Symp., [Pap.], 1st (1974), Meeting Date 1973, 346-62.
Editor(s): Brown, Algie C. Publisher: MEDCOM Press, New York, N. Y.
CODEN: 34QFAU
DT Conference
LA English
AB Human **scalp** and beard **hair follicles**

actively metabolized testosterone-1,2-3H2 in vitro. The principal products formed by both **hair follicle** types after 2 h of incubation were androstanediol, androsterone, dihydrotestosterone, androstenedione, 5 β -androstanedione, 5 α -androstanedione, and a water-insol. ester of dihydrotestosterone. Therefore, there are at least 6 potentially active metabolic pathways for testosterone catabolism in human **scalp** and beard **hair follicles**: (1) the reduction of a 3-one to 3 α -ol; (2) the oxidation of a 17 β -ol to 17-one; (3) the 5 α saturation of a 4-5 double bond; (4) the 5 β saturation of a 4-5 double bond; (5) the esterification of a 17 β -ol; and (6) an unknown pathway, probably also an esterification. Under the conditions of the experiment, testosterone metabolism, testosterone uptake, and total metabolite formation were the same for **scalp** and beard **follicles**. Thus, the enzymic conversion of testosterone to a more powerful androgen may not be significant in the hormonal stimulation of **hair growth**. Similarly, this could also apply to the mol. basis of common **baldness**.

L82 ANSWER 10 OF 11 HCAPLUS COPYRIGHT 2004 ACS on STN
AN 1969:528584 HCAPLUS
DN 71:128584
TI Comparative study of two antidandruff preparations
AU **Orentreich, Norman**; Taylor, Edmund H.; Berger, Robert A.;
Auerbach, Robert
CS Orentreich Med. Group, New York, NY, USA
SO Journal of Pharmaceutical Sciences (1969), 58(10), 1279-80
CODEN: JPMSAE; ISSN: 0022-3549
DT Journal
LA English
AB The relative antidandruff efficacy of a com. available 2% zinc pyrithione shampoo, a 2.5% Se sulfide suspension, and an unmedicated control shampoo was measured using a well-tested visual technique. The zinc pyrithione shampoo and the Se sulfide suspension were equally effective, both being significantly more effective than the control shampoo. A supplemental evaluation of the effects of the test products on **scalp** oiliness is also reported.

L82 ANSWER 11 OF 11 HCAPLUS COPYRIGHT 2004 ACS on STN
AN 1958:7828 HCAPLUS
DN 52:7828
OREF 52:1441d
TI Alkaline phosphatase in **alopecia areata**
AU Kopf, Alfred W.; **Orentreich, Norman**
CS New York Univ., New York, NY
SO AMA Archives of Dermatology (1957), 76, 288-95
CODEN: AMDEAB; ISSN: 0096-5359
DT Journal
LA Unavailable
AB Alkaline phosphatase activity is diminished in hair papillae during early stages of **alopecia areata**. In the intermediate stages the enzyme activity is restored and in the late stages it increases markedly.

=> => fil wpix
FILE 'WPIX' ENTERED AT 16:43:00 ON 19 MAY 2004
COPYRIGHT (C) 2004 THOMSON DERWENT

FILE LAST UPDATED: 14 MAY 2004 <20040514/UP>
MOST RECENT DERWENT UPDATE: 200431 <200431/DW>
DERWENT WORLD PATENTS INDEX SUBSCRIBER FILE, COVERS 1963 TO DATE

>>> FOR A COPY OF THE DERWENT WORLD PATENTS INDEX STN USER GUIDE,

PLEASE VISIT:

http://www.stn-international.de/training_center/patents/stn_guide.pdf <<<

>>> FOR DETAILS OF THE PATENTS COVERED IN CURRENT UPDATES, SEE
<http://thomsonderwent.com/coverage/latestupdates/> <<<

>>> FOR INFORMATION ON ALL DERWENT WORLD PATENTS INDEX USER
 GUIDES, PLEASE VISIT:
<http://thomsonderwent.com/support/userguides/> <<<

>>> NEW! FAST-ALERTING ACCESS TO NEWLY-PUBLISHED PATENT
 DOCUMENTATION NOW AVAILABLE IN DERWENT WORLD PATENTS INDEX
 FIRST VIEW - FILE WPIFV. FREE CONNECT HOUR UNTIL 1 MAY 2004.
 FOR FURTHER DETAILS: <http://www.thomsonderwent.com/dwpifv> <<<

>>> NEW! IMPROVE YOUR LITIGATION CHECKING AND INFRINGEMENT
 MONITORING WITH LITALERT. FIRST ACCESS TO RECORDS OF IP
 LAWSUITS FILED IN THE 94 US DISTRICT COURTS SINCE 1973.
 FOR FURTHER DETAILS:
<http://www.thomsonscientific.com/litalert> <<<

>>> THE DISPLAY LAYOUT HAS BEEN CHANGED TO ACCOMODATE THE
 NEW FORMAT GERMAN PATENT APPLICATION AND PUBLICATION
 NUMBERS. SEE ALSO:
<http://www.stn-international.de/archive/stnews/news0104.pdf> <<<

>>> SINCE THE FILE HAD NOT BEEN UPDATED BETWEEN APRIL 12-16
 THERE WAS NO WEEKLY SDI RUN <<<

=> d all abeq tech abex tot

L116 ANSWER 1 OF 3 WPIX COPYRIGHT 2004 THOMSON DERWENT on STN

AN 2004-034758 [03] WPIX

DNC C2004-011463

TI Reduction of mammalian **hair growth** in cosmetic
 treatment, comprises applying, to area of skin, composition comprising
 thiazolidinone derivative.

DC D21 E13

IN AHLUWALIA, G S; HWANG, C S; JARDIEN, A; SHANDER, D

PA (AHLU-I) AHLUWALIA G S; (HWAN-I) HWANG C S; (JARD-I) JARDIEN A; (SHAN-I)
 SHANDER D; (GILL) GILLETTE CO

CYC 103

PI WO 2003096997 A1 20031127 (200403)* EN 20 A61K007-06

RW: AT BE BG CH CY CZ DE DK EA EE ES FI FR GB GH GM GR HU IE IT KE LS
 LU MC MW MZ NL OA PT RO SD SE SI SK SL SZ TR TZ UG ZM ZW

W: AE AG AL AM AT AU AZ BA BB BG BR BY BZ CA CH CN CO CR CU CZ DE DK
 DM DZ EC EE ES FI GB GD GE GH GM HR HU ID IL IN IS JP KE KG KP KR
 KZ LC LK LR LS LT LU LV MA MD MG MK MN MW MX MZ NI NO NZ OM PH PL
 PT RO RU SC SD SE SG SK SL TJ TM TN TR TT TZ UA UG US UZ VC VN YU
 ZA ZM ZW

US 2003220300 A1 20031127 (200410) A61K031-675

ADT WO 2003096997 A1 WO 2003-US13956 20030502; US 2003220300 A1 US 2002-145283
 20020514

PRAI US 2002-145283 20020514

IC ICM A61K007-06; A61K031-675

ICS A61K008-49; A61K031-426

AB WO2003096997 A UPAB: 20040505

NOVELTY - Reduction of mammalian **hair growth** comprises
 selecting an area of skin from which reduced **hair growth**
 is desired; and applying a composition comprising a thiazolidinone
 derivative in an amount to reduce **hair growth**.

DETAILED DESCRIPTION - INDEPENDENT CLAIMS are also included for:

(1) use of a thiazolidinone derivative in preparation of a medicament

for reducing mammalian **hair growth**; and

(2) a method for the preparation of the composition comprising incorporating a thiazolidinone derivative in a **hair-growth** inhibiting concentration into a carrier.

ACTIVITY - Depilatory.

MECHANISM OF ACTION - None given.

USE - For reducing mammalian **hair growth** (claimed) in cosmetic treatment (claimed). The area of skin may be the face, legs, arms, armpits or torso, the composition being applied in conjunction with shaving. It may be applied to an area of the skin of a woman with hirsutism. The **hair growth** may comprise androgen stimulated **hair growth**.

ADVANTAGE - The composition provides a reduction in **hair growth** of at least 15-35% when tested in the Golden Syrian Hamster assay.

Dwg.0/0

FS CPI

FA AB; GI; DCN

MC CPI: D08-B07; E06-A02E; E07-F01

TECH UPTX: 20040505

TECHNOLOGY FOCUS - ORGANIC CHEMISTRY - Preferred Compounds: The thiazolidinone derivative is of formula (I).

X and Y = oxygen, nitrogen or sulfur;

R1 = aryl;

R2 = H or (CH₂)_nA;

n = 1-5;

A = CO₂H, or SO₃H.

At least X or Y is an oxygen.

The composition may further comprise a second compound that also causes a reduction in **hair growth**. The thiazolidinone derivative has an enantiomeric excess of the active isomer.

Preferred Composition: The thiazolidinone derivative is present in the composition at 0.1-30%. The thiazolidinone derivative is applied to the skin at 10-3000 mug of the compound/cm² of skin.

ABEX UPTX: 20040505

SPECIFIC COMPOUNDS - (I) is **ciglitazone**, **pioglitazone**, **rosiglitazone**, **trogliatzone**, **darglitazone**, **englitazone** or 5-(5-nitro-2-phenylsulfanyl-benzylidene)-2-thioxo-thiazolidin-4-one.

EXAMPLE - A composition contained **pioglitazone** (5 weight%) in a vehicle comprising ethanol (80%) and dimethyl sulfoxide (20%). A 47.7% reduction in hair mass was detected after 3 weeks in the Golden Syrian Hamster assay.

L116 ANSWER 2 OF 3 WPIX COPYRIGHT 2004 THOMSON DERWENT on STN

AN 2003-681209 [65] WPIX

DNC C2003-186257

TI Method for the treatment or prevention of disorders of the skin or exoskeleton, e.g. **hair growth** deficiency, acne or skin cancer, comprises administration of peroxisome proliferator activated receptor ligands, e.g. **rosiglitazone**.

DC B05 D21

IN MOESSNER, R; REICH, K

PA (UYGE-N) UNIV GEORG AUGUST GOETTINGEN

CYC 1

PI DE 10204398 A1 20030814 (200365)* 9 A61K045-00

ADT DE 10204398 A1 DE 2002-10204398 20020204

PRAI DE 2002-10204398 20020204

IC ICM A61K045-00

AB DE 10204398 A UPAB: 20031009

NOVELTY - The use of compounds (I), which interact with at least one peroxisome proliferator activated receptor (PPAR) subtype, for the

production of medicaments for the treatment or prophylaxis of disorders of the skin or exoskeleton.

ACTIVITY - Dermatological; Antialopecia; Antiseborrheic; Antiinflammatory; Cytostatic.

Rosiglitazone (Ia) inhibited the growth of melanoma cells by an average of 76% at a concentration of 20 micro M, and by almost 100% at a concentration of 50 micro M.

MECHANISM OF ACTION - Peroxisome Proliferator Activated Receptor (PPAR) agonist.

USE - Compounds (I) are specifically used for treating or preventing hair growth disorders, diseases of the sebaceous glands (especially acne, rosacea or perioral dermatitis) or malignant tumors of the skin (especially tumors of non-melanocyte epidermal cells, particularly basal cell skin carcinoma, plate epithelial carcinoma of the skin or mucosa or tumors originating in the sebaceous glands of the skin; or malignant melanoma).

ADVANTAGE - Compounds (I) are effective in the treatment or prevention of hair growth disorders, diseases of the sebaceous glands or malignant tumors of the skin.

DESCRIPTION OF DRAWING(S) - The figure shows graphically the quantitative determination of PPAR mRNA in six different melanomas, as determined by quantitative real time RT-PCR.

Dwg.1/6

FS

CPI

FA

AB; GI; DCN

MC

CPI: B04-H03; B07-D04C; B07-F01; B14-H01; B14-L01; B14-N17;

B14-R02; D08-B03; D08-B09A

TECH

UPTX: 20031009

TECHNOLOGY FOCUS - PHARMACEUTICALS - Preferred Components: Compounds (I) are PPAR agonists and/or interact with the gamma-subtype of PPAR. (I) is a natural product, specifically 15-deoxy-DELTA12,14-prostaglandin J2; or a synthetic compound, specifically a thiazolidine dione derivative, especially **rosiglitazone** or **pioglitazone**.

ABEX

UPTX: 20031009

SPECIFIC COMPOUNDS - Three compounds (I) are specifically claimed, i.e. 15-deoxy-DELTA12,14-prostaglandin J2; **Rosiglitazone** (Ia); and **Pioglitazone**.

L116 ANSWER 3 OF 3 WPIX COPYRIGHT 2004 THOMSON DERWENT on STN

AN 2001-565396 [63] WPIX

DNC C2001-167804

TI Use of an insulin sensitivity increasing

substance to treat disorders of the pilosebaceous apparatus e.g. alopecia, acne, hirsutism or superfluous hair growth.

DC B05

IN KRAJCIK, R A; ORENTREICH, N

PA (OREN-N) ORENTREICH FOUND ADVANCEMENT SCI INC

CYC 95

PI WO 2001062237 A2 20010830 (200163)* EN 22 A61K031-00

RW: AT BE CH CY DE DK EA ES FI FR GB GH GM GR IE IT KE LS LU MC MW MZ
NL OA PT SD SE SL SZ TR TZ UG ZW

W: AE AG AL AM AT AU AZ BA BB BG BR BY BZ CA CH CN CR CU CZ DE DK DM
DZ EE ES FI GB GD GE GH GM HR HU ID IL IN IS JP KE KG KP KR KZ LC
LK LR LS LT LU LV MA MD MG MK MN MW MX MZ NO NZ PL PT RO RU SD SE
SG SI SK SL TJ TM TR TT TZ UA UG US UZ VN YU ZA ZW

AU 2001039826 A 20010903 (200202) A61K031-00

US 2002143039 A1 20021003 (200267) A61K031-426

EP 1267850 A2 20030102 (200310) EN A61K031-00

R: AL AT BE CH CY DE DK ES FI FR GB GR IE IT LI LT LU LV MC MK NL PT
RO SE SI TR

ADT WO 2001062237 A2 WO 2001-US5653 20010223; AU 2001039826 A AU 2001-39826
20010223; US 2002143039 A1 Provisional US 2000-184398P 20000223, Cont of

WO 2001-US5653 20010223, US 2002-73607 20020211; EP 1267850 A2 EP 2001-914437 20010223, WO 2001-US5653 20010223

FDT AU 2001039826 A Based on WO 2001062237; EP 1267850 A2 Based on WO 2001062237

PRAI US 2000-184398P 20000223; US 2002-73607 20020211

IC ICM A61K031-00; A61K031-426
ICS A61K031-155

AB WO 200162237 A UPAB: 20011031

NOVELTY - Use of an **insulin sensitivity increasing substance (ISIS)** for treating disorders of the **pilosebaceous** apparatus is new.

DETAILED DESCRIPTION - INDEPENDENT CLAIMS are also for the following:

- (a) Use of the **ISIS** in combination with a steroid enzyme inhibitor or inducer (STI);
- (b) Use of the **ISIS** in combination with an androgen receptor blocking agent (ARB);
- (c) Use of the **ISIS** in combination with an activity-enhancing agent and optionally an STI or ARB; and
- (d) A composition comprising an **ISIS** in combination with at least one ARB and an STI.

ACTIVITY - Antiseborrheic; Dermatological; Depilatory.

MECHANISM OF ACTION - Steroid Enzyme Inhibitor; Androgen Receptor Blocking Agent.

USE - The compositions are used to treat disorders of the **pilosebaceous** apparatus (**hair/oil gland**) including inhibiting, reducing or reversing **hair loss**. The compositions may be used to treat **alopecia**, acne, hirsutism or superfluous **hair growth**.

Dwg.0/0

FS CPI

FA AB; DCN

MC CPI: B01-C04; B06-A01; B07-D04C; B07-D09; B07-F01; B10-A17; B10-E04A

TECH UPTX: 20011031

TECHNOLOGY FOCUS - PHARMACEUTICALS - Preferred **Insulin Sensitivity: Increasing Substance: ISISs** include a biguanidine, metoformin hydrochloride, a **thiazolidinedione** selected from **troglitazone**, **ciglitazone**, **piolitazone**, **rosiglitazone** and **englitazone** and **d-chiro-inositol**.

Preferred Steroid Enzyme Inhibitor or Inducer: A 5-alpha reductase inhibitor, a 3-alpha hydroxy steroid dehydrogenase inhibitor or a 17-beta hydroxy steroid hydrogenase inducer.

Preferred Androgen Receptor Blocking Agent; Cytoproterone acetate, flutamide, bicalutamide, nilutamide, RU-58841, canrenone, spironolactone, progesterone, 4mA, ketoconazole and cimetidine. Preferred Activity Enhancing Agent: A penetration enhancing agent, a vasodilator, an anti-inflammatory, a glucose/**insulin** regulating compound or an endogenous or exogenous effector.

ABEX UPTX: 20011031

ADMINISTRATION - Administration may be oral, by injection or preferably topical (claimed). No dosage is given.

EXAMPLE - The effects of **metformin** administration on **hair loss** in aged female obese mice was studied.

Metformin HCL was dissolved in drinking water at a dosage of 240 mg/kg/d. The dosage is adjusted weekly to reflect changes in body weight and the water changed twice weekly. Ob/ob (homozygous) and ob/+ heterozygous mice were used. The ob/+ mice were not entered into the study until 2 weeks after the ob/ob mice as no mice were available at the age group required for the studies (7 weeks). The mice were provided with food and drink ad libitum and their body weight and food and water consumption monitored twice a week. Blood samples were also collected to monitor serum glucose and insulin levels. The results of the study

indicated that onset of **hair loss** in female ob/ob mice is delayed by **metformin** treatment. Clear differences between control and **metformin** treated animals were detected by 10 months of age (corresponding to nearly 8 months of treatment). At that time, **hair loss** was observed in 36% of the control ob/ob mice, while only 8% of the **metformin** treated animals were affected. **Hair loss** was not observed in control or **metformin** treated ob/+ mice.

=> d his

(FILE 'HOME' ENTERED AT 15:44:02 ON 19 MAY 2004)
SET COST OFF

FILE 'REGISTRY' ENTERED AT 15:44:21 ON 19 MAY 2004

	E THIAZOLIDINEDIONE/CN
L1	1 S E3
	E TROGLITAZONE/CN
L2	1 S E3
	E CIGLITAZONE/CN
L3	1 S E3
	E PIOGLITAZONE/CN
L4	1 S E3
	E ROSIGLITAZONE/CN
L5	1 S E3
	E ENGLITAZONE/CN
L6	1 S E3
	E D-CHIRO-INOSITOL/CN
L7	1 S E3
	E METFORMIN/CN
L8	2 S E3, E5
	E CYTOPROTERONE/CN
	E CITOPROTERONE/CN
	E CYPROTERONE/CN
L9	2 S E3-E6
	E FLUTAMIDE/CN
L10	1 S E3
	E BICALUTAMIDE/CN
L11	1 S E3
	E NILUTAMIDE/CN
L12	1 S E3
	E RU-58841/CN
	E RU 58841/CN
L13	1 S E3
	E CANRENONE/CN
L14	1 S E3
	E SPIRONOLACTONE/CN
L15	1 S E3
	E PROGESTERONE/CN
L16	1 S E3
	E 4MA/CN
	E 4 MA/CN
	E 4-MA/CN
L17	1 S E3
	E KETOCONAZOLE/CN
L18	1 S E3
	E CIMETIDINE/CN
L19	1 S E3
L20	9 S L1-L8
	SEL RN
L21	227 S E1-E9/CRN
L22	217 S L21 NOT MXS/CI

L23 62 S L22 NOT (COMPD OR WITH)
L24 57 S L23 NOT IDS/CI
L25 12 S L9-L19
SEL RN
L26 284 S E10-E21/CRN
L27 17 S L26 NOT ((MXS OR IDS)/CI OR COMPD OR WITH OR UNSPECIFIED)
L28 15 S L27 NOT CONJUGATE

FILE 'HCAPLUS' ENTERED AT 15:58:46 ON 19 MAY 2004

L29 4690 S L20 OR L24
L30 2291 S D() (CHIROINOSITOL OR CHIRO INOSITOL) OR METFORMIN# OR DIMETHY
L31 2672 S ROSIGLITAZON# OR BRL49653 OR BRL() (49653 OR 49 653) OR PIOGLI
L32 2939 S CS045 OR CS 045 OR GR92132# OR GR() (92132# OR 92 132#) OR ADD
L33 6815 S L29-L32
L34 1970 S ISIS OR INSULIN(L) SENSITIV? (L) INCREAS? (L) SUBSTANC?
L35 12995 S ALOPEC? OR BALDNESS OR BALD OR BALDING OR HAIR(L) (LOSS OR LOS
L36 170 S PILOSEBAC?
L37 3598 S HAIR(L)?FOLLIC?
L38 3520 S SCALP?
L39 14 S L33 AND L35-L38
L40 3 S L34 AND L35-L38
L41 16 S L39,L40
E HAIR/CT
L42 16601 S E3-E18
E E3+ALL
L43 30975 S E6,E5+NT
E E15+ALL
L44 2450 S E13,E12+NT
E E15+ALL
E E17+ALL
L45 20263 S E2+NT
E E8+ALL
E E19+ALL
L46 2865 S E7,E6+NT
E E16+ALL
E E20+ALL
L47 871 S E4
E E6+ALL
E E21+ALL
L48 2215 S E3,E2+NT
E HAIR/CT
L49 65 S E86-E88
L50 15265 S E44-E68
L51 32 S L33 AND L42-L50
L52 1 S L34 AND L42-L50
L53 35 S L41,L51,L52
L54 55669 S L25 OR L28
L55 82367 S CYPROTERON? OR CYPROTERON?(S) ACETATE OR FLUTAMID? OR BICALUTA
L56 213 S L55 AND L33
L57 6 S L55 AND L34
L58 14 S L56,L57 AND L53
L59 764 S 17 BETA HYDROXY STEROID DEHYDROGENASE
L60 1313 S 17 BETA HYDROXYSTEROID DEHYDROGENASE
L61 320 S 3 ALPHA HYDROXY STEROID DEHYDROGENASE
L62 736 S 3 ALPHA HYDROXYSTEROID DEHYDROGENASE
L63 3272 S 5 ALPHA REDUCTASE

FILE 'REGISTRY' ENTERED AT 16:13:59 ON 19 MAY 2004

L64 3 S 9015-81-0 OR 9028-56-2 OR 9081-34-9

FILE 'HCAPLUS' ENTERED AT 16:14:07 ON 19 MAY 2004

L65 5023 S L64
L66 6 S L33 AND L59-L63,L65

L67 1 S L34 AND L59-L63,L65
 L68 6 S L66,L67
 L69 19 S L58,L68
 L70 4 S L69 AND ALOPEC?
 L71 4 S L35 AND L69
 L72 4 S L70,L71
 L73 175 S L54 AND L33,L34
 L74 3 S L73 AND L35-L38
 L75 11 S L73 AND L42-L50
 L76 19 S L69-L72,L74,L75
 E KRAJCIK R/AU
 L77 8 S E4,E6,E7
 E ORENTREICH N/AU
 L78 45 S E3,E4
 L79 1 S US20020143039/PN OR (WO2001-US5653 OR US2000-184398#)/AP,PRN
 L80 12 S L77-L79 AND L29-L63,L65
 L81 1 S L80 AND L76
 L82 11 S L80 NOT L81
 L83 18 S L76 NOT L81
 L84 6 S L83 AND (PD<=20000223 OR PRD<=20000223 OR AD<=20000223)
 L85 12 S L83 NOT L84
 SEL DN AN 1
 L86 1 S L85 AND E1-E3
 L87 2 S L79,L86
 L88 2 S L87 AND L1-L28
 L89 2 S L88 AND L29-L63,L65-L88

FILE 'REGISTRY' ENTERED AT 16:28:47 ON 19 MAY 2004

L90 1 S 56-03-1
 L91 1 S 102-02-3

FILE 'HCAPLUS' ENTERED AT 16:28:58 ON 19 MAY 2004

L92 1269 S L90 OR L91
 L93 4 S L92 AND L35-L38
 L94 8 S L92 AND L42-L50
 L95 8 S L93,L94
 SEL DN AN 3-7
 L96 5 S L95 AND E4-E18
 L97 6 S L89,L96

FILE 'HCAPLUS' ENTERED AT 16:31:26 ON 19 MAY 2004

FILE 'WPIX' ENTERED AT 16:32:35 ON 19 MAY 2004

L98 945 S L30/BIX OR L31/BIX OR L32/BIX
 L99 104 S L34/BIX
 E THIAZOLIDINEDIONE/DCN
 E THIAZOLIDINE/DCN
 E E7+ALL
 L100 24 S E2
 E THIAZOLIDINEDIONE/CN
 L101 1 S E2
 E TROGLITAZONE/DCN
 E TROGLITAZONE/CN
 L102 3 S E3-E5
 E REZULIN/DCN
 E CIGLITAZONE/DCN
 E PIOGLITAZONE/DCN
 E E3+ALL
 L103 213 S E2
 E ROSIGLITAZONE/DCN
 E ENGLITAZONE/DCN
 E D-CHIRO-INOSITOL/DCN
 E CHIRO-INOSITOL/DCN

L104 E E2+ALL
 42 S E2
 E BIGUANIDE/DCN
 E E3+ALL
L105 98 S E2
 E METFORMIN/DCN
 E E3+ALL
L106 253 S E2
L107 1192 S L98-L106
L108 16 S L107 AND L35/BIX
L109 2 S L107 AND A61P017-14/IC, ICM, ICS, ICA, ICI
L110 0 S L107 AND A61P017:14/ICI
L111 15 S L107 AND (B14-R02 OR C14-R02 OR B12-L05 OR C12-L05 OR D08-B03
L112 24 S L108-L111
L113 1 S L107 AND (L36/BIX OR L37/BIX OR L38/BIX)
L114 24 S L112, L113
L115 24 S L114 AND L98-L114
 SEL DN AN 5 7 16
L116 3 S L115 AND E1-E6

FILE 'WPIX' ENTERED AT 16:43:00 ON 19 MAY 2004

=>